

Linear Congruence Equations

Let $a, x \in \mathbf{Z}_n^*$. Recall that x is said to be an *inverse* of a modulo n if $ax \equiv 1 \pmod{n}$. It is easily seen that the inverse, if it exists, is unique modulo n , for if $ax \equiv 1 \pmod{n}$ and $ay \equiv 1 \pmod{n}$, then $x \equiv xay \equiv y \pmod{n}$. We denote this unique x , when it exists, by $a^{-1} \pmod{n}$ (or simply a^{-1} when the modulus n is clear from context).

Theorem 1 *Let $a \in \mathbf{Z}_n^*$. Then a^{-1} exists in \mathbf{Z}_n^* .*

Proof: Let $a \in \mathbf{Z}_n^*$ and consider the function $f_a(x) = ax \pmod{n}$. f_a is easily shown to be a one-one mapping from \mathbf{Z}_n^* to \mathbf{Z}_n^* . Hence, f_a is also onto, so for some $x \in \mathbf{Z}_n^*$, $f_a(x) = 1$. Then $ax \equiv 1 \pmod{n}$, so $x = a^{-1} \pmod{n}$. ■

We showed in class how to use the Extended Euclidian algorithm to efficiently compute $a^{-1} \pmod{n}$ given a and n .

Here we consider the solvability of the more general linear congruence equation:

$$ax \equiv b \pmod{n} \tag{1}$$

where $a, b \in \mathbf{Z}_n^*$ are constants, and x is a variable ranging over \mathbf{Z}_n^* .

Theorem 2 *Let $a, b, n \in \mathbf{Z}_n^*$. Let $d = \gcd(a, n)$. If $d \mid b$ then $ax \equiv b \pmod{n}$ has d solutions x_0, \dots, x_{d-1} , where*

$$x_t = \left(\frac{b}{d}\right)\bar{x} + \left(\frac{n}{d}\right)t \tag{2}$$

and $\bar{x} = \left(\frac{a}{d}\right)^{-1} \pmod{\left(\frac{n}{d}\right)}$. If $d \nmid b$, then $ax \equiv b \pmod{n}$ has no solutions.

Proof: Let $d = \gcd(a, n)$. Clearly if $ax \equiv b \pmod{n}$, then $d \mid b$, so there are no solutions if $d \nmid b$.

Now suppose $d \mid b$. Since $\left(\frac{a}{d}\right)$ and $\left(\frac{n}{d}\right)$ are relatively prime, \bar{x} exists by Theorem 1. Multiplying both sides of (2) by a , we get

$$ax_t = b \left(\frac{a}{d}\right)\bar{x} + n \left(\frac{a}{d}\right)t \tag{3}$$

where now we are working over the integers. But $\left(\frac{a}{d}\right)\bar{x} = 1 + \frac{kn}{d}$ for some k by the definition of \bar{x} , so substituting for $\left(\frac{a}{d}\right)\bar{x}$ in (3) yields

$$ax_t = b + kn \left(\frac{b}{d}\right) + n \left(\frac{a}{d}\right)t \tag{4}$$

The quantities in parentheses are both integers, so it follows immediately that $ax_t \equiv b \pmod{n}$ and hence x_t is a solution of (1).

It remains to show that the d solutions above are distinct modulo n . But this is obvious since $x_0 < x_1 < \dots < x_{d-1}$ and $x_{d-1} - x_0 = \frac{n}{d}(d-1) < n$. ■